

DUKE NEWS RELEASE FORM (paste this at top of your release)

Author: Abby Olena

Directory/Slug: Shared > Releases > Leadpb

Edited by: KArlos

Faculty reviewed: 6 January 2019

Proofed by: RobinAnn

Photo – Shared > Releases > P factor.jpg

Caption - Higher blood lead levels at age 11 are linked to an increase in symptoms of psychopathology and difficult personality traits by age 38.

Photo credit- Adapted from JAMA Psychiatry, 2019;76(4):1-9.

Did you include name of funder in story? Y

DISTRIBUTION /POSTING:

Distribution date (including year), time: Thursday, Jan. 17, 2019

Embargo date, time: January 23, 2019, 11 am EST

Did you post this online?

If yes, did you include the URL at the end of the release?

Vocus contact lists:

Vocus outlet lists (which desk?):

Headline for Web, Eurekalert, Email Subject (Keep it snappy; no more than 7 words; Capitalize First Letter In Each Word): Childhood Lead Exposure Linked to Poor Adult Mental Health

Subhead (Sentence case): Lead exposure in childhood appears to have long-lasting negative effects on mental health and personality in adulthood.

EurekaAlert:

post date -

EurekaAlert 75-word Teaser (Can just be lead paragraphs): Lead exposure in childhood appears to have long-lasting negative effects on mental health and personality in adulthood, according to a study of people who grew up in the era of leaded gasoline. The findings, which will appear Jan. 23 in JAMA Psychiatry, reveal that the higher a person's blood lead levels at age 11, the more likely they are to show signs of mental illness and difficult personality traits by age 38.

Primary Keyword - Medicine/Health

keywords - pollution/remediation; neurochemistry; environmental health;

neurobiology; pediatrics; toxicology; mental health; depression/anger

DOI Number – 10.1001/jamapsychiatry.2018.4192

Funders – New Zealand Health Research Council, New Zealand Ministry of Business, Innovation, and Employment, National Institute on Aging, National Institute of Child Health and Human Development, National Institute of Environmental Health Sciences, UK Medical Research Council, The Jacobs Foundation, The Avielle Foundation

Grant Numbers: T32AG000139, AG032282, T32HD007376, F31ES029358,
MR/P005918/1

EurekAlert Tweet (150 characters): Children exposed to lead may have poor mental health as adults, according to a study of people born in the 1970s.

EurekAlert Multimedia (repeat as necessary):

Title - Childhood Lead Exposure Tied to Poor Adult Mental Health

Caption - Higher blood lead levels at age 11 are linked to an increase in symptoms of psychopathology and difficult personality traits by age 38.

Credit - Adapted from JAMA Psychiatry, 2019;76(4):1-9.

Permissions (all or registered reporters)

Usage restrictions (if any)

Duke Today Topics/Tagwords: Psychology and Neuroscience, Psychiatry & Behavioral Sciences, Graduate School, Environment, Research

Duke Tweet (120 characters): Children exposed to lead may have poor mental health as adults.

DUKE UNIVERSITY NEWS
Office of News & Communications
<http://today.duke.edu>

EMBARGOED FOR RELEASE: 11 am EST, Wednesday, January, 23, 2019

CONTACT: Karl Bates
(919) 681-8054
karl.bates@duke.edu

CHILDHOOD LEAD EXPOSURE LINKED TO POOR ADULT MENTAL HEALTH

DURHAM, N.C. -- Lead exposure in childhood appears to have long-lasting negative effects on mental health and personality in adulthood, according to a study of people who grew up in the era of leaded gasoline.

Previous studies have identified a link between lead and intelligence, but this study looked at changes in personality and mental health as a result of exposure to the heavy metal.

The findings, which will appear Jan. 23 in *JAMA Psychiatry*, reveal that the higher a person's blood lead levels at age 11, the more likely they are to show signs of mental illness and difficult personality traits by age 38.

The link between mental health and lead exposure is modest, according to study coauthor Aaron Reuben, a graduate student in clinical psychology at Duke University. But "it's potentially important because this is a modifiable risk factor that at one point in time everyone was exposed to, and now, certain people in certain cities and countries are still exposed to," he said.

In a previous study, Reuben and colleagues showed that higher levels of lead in childhood were linked to lower IQ and lower social standing in adulthood.

Both sets of findings suggest that lead's "effects really can last for quite a long time, in this case three to four decades," said coauthor Jonathan Schaefer, also a graduate student in clinical psychology at Duke. "Lead exposure decades ago may be harming the mental health of people today who are in their 40s and 50s."

Because gasoline around the world was treated with high levels of lead from the mid 1960s until the late 1980s, most adults now in their 30s, 40s, and 50s were exposed as children. Lead from automotive exhaust was released into the atmosphere and soils. Today, high lead exposures are rarer, and most often found in children who live in older buildings with lead plumbing and paint.

The subjects of this study are part of a group of more than 1,000 people born in 1972 and 1973 in Dunedin, New Zealand, at a time when gasoline lead levels in New Zealand were among the highest in the world. They have regularly participated in physical and mental health evaluations at the local University of Otago.

Researchers measured blood lead levels -- in micrograms per deciliter of blood (ug/dL) -- when participants were 11 years old. Today, blood lead levels above 5 ug/dL will trigger additional clinical follow-up of a child. At age 11, 94 percent of participants in the Dunedin Study had blood lead levels above this cutoff.

"These are historical data from an era when lead levels like these were viewed as normal in children and not dangerous, so most of our study participants were never given any treatment for lead toxicity," said Terrie Moffitt, the senior author of the study and Duke's Nannerl O. Keohane University Professor of psychology & neuroscience and psychiatry & behavioral sciences.

The Duke research team also assessed participant mental health and personality at various points throughout their lives, most recently at age 38. Diagnostic criteria or symptoms associated with eleven different psychiatric disorders -- dependence on alcohol, cannabis, tobacco, or hard drugs; conduct disorder, major depression, generalized anxiety disorder, fears and phobias, obsessive-compulsive disorder, mania, and schizophrenia -- were used to calculate a single measure of mental health, called the psychopathology factor, or "p-factor" for short.

The higher an individual's p-factor score, the greater the number and severity of psychiatric symptoms. Lead's effects on mental health as measured by the p-factor score are about as strong as those on IQ, explained coauthor Avshalom Caspi, Edward M. Arnett Professor of psychology & neuroscience and psychiatry & behavioral sciences at Duke. "If you're worried about lead exposure's impact on IQ, our study suggests you should probably also be worried about mental health," Caspi said.

The research team also determined that participants exposed to higher levels of lead as children were described as having more difficult adult personalities by family members and friends. Specifically, they found that study members with greater lead exposure were rated as more neurotic, less agreeable, and less conscientious than their less-exposed peers.

These findings confirm personality characteristics that have been previously linked to a number of problems, including worse mental and physical health, reduced job satisfaction, and troubled interpersonal relationships.

"For folks who are interested in intervention and prevention, the study suggests that if you're going to intervene on a group of kids or young adults that have been lead exposed, you may need to think very long-term when it comes to their care," said Schaefer.

In the future, the Dunedin Study team is interested in whether lead exposure might be linked to the development of later-life diseases such as dementia or cardiovascular disease.

Reuben said the findings are relevant to other developed countries as well. "When we see changes that may be the result of lead exposures in New Zealand it's very likely that you would have seen those same impacts in America, in Europe, and the other countries that were using leaded gasoline at the same levels at the same time."

The New Zealand Health Research Council and the New Zealand Ministry of Business, Innovation, and Employment provided funding to the Dunedin Multidisciplinary Health and

Development Research Unit. Support also came from grants T32AG000139 and AG032282 from the National Institute on Aging, T32HD007376 from the National Institute of Child Health and Human Development, F31ES029358 from the National Institute of Environmental Health Sciences, and MR/P005918/1 from the UK Medical Research Council. The Jacobs Foundation and the Avielle Foundation provided additional funding.

CITATION: "Association of Childhood Lead Exposure With Adult Personality Traits and Lifelong Mental Health," Aaron Reuben, Jonathan D. Schaefer, Terrie E. Moffitt, Jonathan Broadbent, Honalee Harrington, Renate M. Houts, Sandhya Ramrakha, Richie Poulton, Avshalom Caspi. JAMA Psychiatry, January 23, 2019.

DOI:10.1001/jamapsychiatry.2018.4192

<https://jama.jamanetwork.com/article.aspx?doi=10.1001/jamapsychiatry.2018.4192>

#