

# MAOA, Childhood Maltreatment, and Antisocial Behavior: Meta-analysis of a Gene-Environment Interaction

Amy L. Byrd and Stephen B. Manuck

**Background:** In a seminal study of gene-environment interaction, childhood maltreatment predicted antisocial behavior more strongly in male subjects carrying an *MAOA* promoter variant of lesser, compared with higher, transcriptional efficiency. Many further investigations have been reported, including studies of other early environmental exposures and female subjects. Here, we report a meta-analysis of studies testing the interaction of *MAOA* genotype and childhood adversities on antisocial outcomes in predominantly nonclinical samples.

**Methods:** Included were 27 peer-reviewed, English-language studies published through August, 2012, that contained indicators of maltreatment or other family (e.g., parenting, sociodemographic) hardships; *MAOA* genotype; indices of aggressive and antisocial behavior; and statistical test of genotype-environment interaction. Studies of forensic and exclusively clinical samples, clinical cohorts lacking proportionally matched control subjects, or outcomes nonspecific for antisocial behavior were excluded. The Liptak-Stouffer weighted Z-test for meta-analysis was implemented to maximize study inclusion and calculated separately for male and female cohorts.

**Results:** Across 20 male cohorts, early adversity presaged antisocial outcomes more strongly for low-activity, relative to high- activity, *MAOA* genotype ( $p = .0044$ ). Stratified analyses showed the interaction specific to maltreatment ( $p = .0000082$ ) and robust to several sensitivity analyses. Across 11 female cohorts, *MAOA* did not interact with combined early life adversities, whereas maltreatment alone predicted antisocial behaviors preferentially, but weakly, in female subjects of high-activity *MAOA* genotype ( $p = .02$ ).

**Conclusions:** We found common regulatory variation in *MAOA* to moderate effects of childhood maltreatment on male antisocial behaviors, confirming a sentinel finding in research on gene-environment interaction. An analogous, but less consistent, finding in female subjects warrants further investigation.

---

**Key Words:** Antisocial behavior, childhood maltreatment, gene-environment interaction, genetics, *MAOA*, meta-analysis

---

Two widely cited reports of putative gene-environment ( $G \times E$ ) interaction, published in 2002 and 2003, described genotype-dependent environmental influences on risk for antisocial behavior and depression, respectively, in a well-characterized, longitudinally studied birth cohort (1,2). In the second of these studies, recent stressful life events and childhood maltreatment predicted depression in young adults in proportion to the number of short (deletion) alleles carried of a 44-base pair insertion/deletion (long/short) polymorphism in the regulatory region of the serotonin transporter gene (serotonin transporter gene-linked polymorphic region [5-HTTLPR]) (2). Two prominent meta-analyses published in 2009 failed to confirm replication of this most frequently cited instance of gene-stress interaction (3,4). Some authors cautioned that these analyses incorporated only a fraction of relevant investigations and relied disproportionately on studies of self-reported life events, rather than contextually sensitive interviews or objective indicators of stress (5,6). In a more comprehensive meta-analysis published subsequently, Karg *et al.* (7) found the 5-HTTLPR to moderate effects of adversity on clinical depression and depressive symptomatology across all published studies ( $p = .00002$ ) but with

some variation among studies of differing methodology. In stratified analyses, 5-HTTLPR genotype interacted with self-reported life events only marginally in predicting depression, whereas the interaction proved robust for studies of childhood maltreatment and of cohorts uniformly exposed to the same stressor or where life events were assessed by structured interview.

The purpose of this article is to extend meta-analytic review to the first of the two  $G \times E$  interactions described by Caspi *et al.* (1). There, exposure to maltreatment in childhood predicted later aggressive and antisocial behaviors among male subjects as a function of regulatory variation in the gene encoding monoamine oxidase-A (*MAOA*). A degradative enzyme, *MAOA* preferentially deaminates the neurotransmitters, serotonin and norepinephrine, and the *MAOA* gene contains a 30-base pair repeating sequence (variable number of tandem repeats [VNTR]) in the 5'-flanking region conferring allele-specific variation in *MAOA* promoter activity (8–10). In this study, early indicators of maltreatment, such as boys' physical or sexual abuse, maternal rejection, or harsh physical punishment, more strongly predicted later conduct problems, antisocial disposition, and violent offending among persons carrying the *MAOA* repeat variant (allele) of lesser transcriptional efficiency (low-activity *MAOA* genotype) than in those of an alternate (high-activity) genotype. This finding has since been cited over 2800 times and prompted similar studies by other investigators. In 2007, Taylor and Kim-Cohen (11) confirmed the interaction of early maltreatment and *MAOA* genotype on antisocial outcomes by meta-analysis of the original study and seven attempted replications (12–18). These studies all included male participants recruited from largely normal populations (*viz.*, excluding forensic or predominantly clinical samples) and contained either a single or composite index of antisocial behavior. Studies also included a measure of participants' childhood exposure to abuse, neglect, or other harm within the family environment, and all

---

From the Department of Psychology, University of Pittsburgh, Pittsburgh, Pennsylvania.

Address correspondence to Amy L. Byrd, M.S., University of Pittsburgh, Department of Psychology, Sterling Plaza, Suite 408, 201 N Craig St, Pittsburgh, PA 15213; E-mail: [alb202@pitt.edu](mailto:alb202@pitt.edu).

Received Jan 22, 2013; revised Apr 20, 2013; accepted May 3, 2013.

reported such exposure positively associated with study outcomes. Consistent with the initial report of Caspi *et al.* (1), the pooled estimate of correlation between family adversity and indices of later antisocial behavior was greater in individuals of low-activity, compared with high-activity, *MAOA* genotype ( $p < .0001$ ) (11).

Many additional investigations have been reported since publication of Taylor and Kim-Cohen (11). These include further G×E studies of early maltreatment (19–25), studies of other environmental moderators (e.g., neighborhood and family socioeconomic disadvantage, peer deviance, parenting styles, commonly experienced life events, maternal prenatal smoking [25–32]), and studies including female subjects (20,21,23, 29–31,33–38) or primarily nonwhite samples (24,33,34). Here, we report a further meta-analysis of the accumulated literature addressing interactions of *MAOA* variation and environmental risk factors in the prediction of aggressive and antisocial outcomes. To allow comparison with the parallel literature on 5-HTTLPR variation, life stress, and depression [i.e., the second G×E literature emerging from the two seminal studies of Caspi *et al.* (1,2)], we have followed the same analytic procedures employed by Karg *et al.* (7). This approach maximizes inclusion of studies of differing design and analytic strategy or of limited statistical reporting or data availability through application of the common Liptak-Stouffer weighted Z-test for meta-analysis, which combines published reports by tests of statistical significance (7,39–42).

## Methods and Materials

### Studies

We sought all peer-reviewed, English-language studies published through August 2012 from: 1) reference lists of prior meta-analysis, narrative review, and individual studies; and 2) major publication databases (e.g., PubMed), using as keywords monoamine oxidase-A or *MAOA* and childhood maltreatment, abuse, adversity or family/family environment and antisocial behavior, conduct disorder/problems, delinquency, externalizing behavior, aggression, or violence. We only included studies that had genotyped the same *MAOA* VNTR reported by Caspi *et al.* (1), those in which the interaction of *MAOA* genotype and early adversity was tested explicitly; and studies for which outcomes included behaviors or disorders on an externalizing or antisocial spectrum (but not solely alcohol or substance abuse). Following Taylor and Kim-Cohen (11), studies of forensic populations (43–45), exclusively clinical samples or clinical samples lacking proportionally matched control subjects, were excluded (46,47). Also excluded were two studies in which indicators of antisocial behavior could not be distinguished from other life outcomes or events (e.g., financial losses, accidents, unspecified relationship problems, or socioeconomic attainments) (48,49).

We identified 27 independent investigations meeting the foregoing criteria and totaling >18,400 study participants. Of these, 12 studies included only male participants (12–16,19,22,24,25,27,28,32), 11 included participants of both sexes (1,17,18,20,21,23,29–31,33,34), and 4 included only female subjects (26,36–38). Study samples were mainly all white (23), with four studies of mixed ethnicity or primarily nonwhite samples (17,19,24,33). To determine if outcomes might vary by similarity to the first reported G×E interaction for *MAOA*, we stratified investigations into two groups: studies focusing specifically on early maltreatment and studies of other childhood adversities. Assignment to the maltreatment group was made when factors such as physical or sexual abuse, assault, or other victimization; severe physical punishment; other exposures to violence, neglect, or

court-mediated family interventions predominated in indices of early childhood environment. Studies of sociodemographic variables, peer affiliations, maternal prenatal smoking, general life events, or parenting styles and those with only minor or oblique representation of maltreatment indicators were assigned to the category of other childhood adversities. When studies reported on measurements from both categories, these were treated as independent tests in stratified analyses.

### Extraction of *p* Values

The authors independently extracted *p* values from each study, without discrepancy. When nonsignificant findings were reported without exact *p* values, we requested more precise values from study authors. If these data were not available or authors declined, we assigned a *p* value of 1 (indicating absence of an interaction implicating either low- or high-activity *MAOA* genotype). Because interaction terms were occasionally collapsed over sex or ethnicity, we also requested *p* values from study authors for male and female subjects analyzed separately and similarly for white and nonwhite segments of multiethnic samples. With respect to outcome measures, for primary analyses, we used the *p* value associated with the most general (e.g., composite) measure of antisocial behavior or computed a weighted mean *p* value when multiple dependent variables and/or multiple environmental moderators were analyzed separately in the published report.

### Genotypes of *MAOA*

Because *MAOA* is located on the X chromosome (Xp11.4-Xp11.3), all male subjects are hemizygous for a single allele of the upstream VNTR, for which variants of 2, 3, 3.5, 4, and 5 repeats have been described. In samples of European ancestry, the 3- and 4-repeat alleles account for >95% of variation. The 2, 3, and 5 repeat variants are commonly grouped as low-activity alleles and contrasted with high-activity alleles of 3.5 or 4 repeats, based on *in vitro* studies of *MAOA* promoter activity (8,10). Functional characterization of the 5 repeat is somewhat controversial (10,20), however, though its rarity suggests a negligible effect on study outcomes resulting from differences in allele grouping. Also, some studies disregarded the rare variants altogether, analyzing only the 3- and 4-repeat alleles.

In female subjects, uncertainty regarding extent of X-inactivation at the *MAOA* locus (50,51) has occasioned differing analytic strategies for comparing *MAOA* genotypes and in some instances served as rationale for excluding female subjects from study analyses (15). When tested in female subjects, some investigators compared only individuals homozygous for low- or high-activity variants (17,21,33,34,37), whereas others included a heterozygous grouping defined by presence of both a low- and high-activity allele (18,20,23,26,29–31,36,38). Here, we treat *MAOA* genotypes in female subjects as they were operationalized in each study; although when a *p* value for the contrast of homozygous low- and high-activity participants was available in studies including heterozygotes, we used this value in the meta-analysis to enhance comparability among studies. Finally, because the sentinel report by Caspi *et al.* (1) addressed risk for antisocial behavior specifically in male subjects and owing to the more variable classification of *MAOA* genotypes in studies of female subjects, analyses were conducted separately for each sex.

### Statistical Analysis

Like the Karg *et al.* (7) meta-analysis of 5-HTTLPR variation, stress, and depression, we combined investigations by the Liptak-Stouffer Z score procedure to yield an aggregate outcome based

on significance tests from each study, adjusted for sample size. Extracted  $p$  values were first expressed as one-tailed metrics, where  $p$  values less than .50 corresponded to liability for antisocial behavior associated with low-activity MAOA genotype and a  $p$  value greater than .50 corresponded to liability for antisocial behavior associated with high-activity MAOA genotype. More precisely, a study outcome was considered consistent with Caspi *et al.* (1) when the dependent variable associated more strongly with an environmental risk factor among participants of low-activity, relative to high-activity, MAOA genotype. As in Karg *et al.* (7),  $p$  values were next converted to Z scores, to which positive and negative signs were attached, respectively, for  $p$  values less than and greater than .50. Finally, a composite Z score was calculated by the formula:

$$Z_w = \frac{\sum_{i=1}^k W_i Z_i}{\sqrt{\sum_{i=1}^k W_i^2}}$$

where  $Z_i$  denotes Z scores of the individual studies,  $W_i$  refers to the study sample size, and  $k$  is the number of studies included in the analysis. The outcome,  $Z_w$ , is then tested for two-tailed significance by reference to the standard normal distribution.

This statistic was first calculated for all studies together and then, in stratified analyses, for studies partitioned by category of environmental moderator (i.e., maltreatment; other adversities). As noted, these analyses were run separately by sex. In four samples that included both male and female subjects, sex-specific  $p$  values were unavailable (17,33,34). We excluded these investigations in primary analyses, but because male subjects comprised a majority of participants in each, we included these with all other male studies in a secondary analysis.

Any initially significant finding was probed for disproportionate influence of single investigations by recomputing  $Z_w$  after removing each study individually, and if found unaltered by deletion of individual studies, the following additional sensitivity analyses were conducted. As noted previously, the  $p$  value entered into analysis was occasionally averaged over tests involving more than one environmental moderator or, more commonly, over multiple outcomes, such as clinical diagnoses, forensic status (e.g., criminal convictions), and dimensional measures of informant and self-reported aggressive, antisocial, or other externalizing behavior. To determine if results were robust to this variation, we repeated the analysis by: 1) iteratively substituting individual outcome variables for composite measures or averaged  $p$  values; and 2) running 1000 additional iterations of the meta-analysis, in each of which a single dependent measure was selected randomly from all studies with multiple outcomes. We then also ran analyses separately for: 1) continuous and dichotomous measures of antisocial behavior; 2) outcomes occurring in childhood/adolescence ( $\leq 18$ ) and those of adulthood; 3) outcomes of overt aggression or violence, nonviolent antisocial behaviors (e.g., vandalism, theft), and measures combining violent and nonviolent indicators (e.g., conduct and antisocial personality disorder diagnoses or symptom counts); and 4) investigations based on cross-sectional and longitudinally studied cohorts. Regarding maltreatment studies, we also conducted analyses separately for those in which environmental exposures were assessed by family (self or parent) report only and studies including nonfamilial informant sources (e.g., official record, observation). Finally, to gauge potential publication bias, we: 1) computed a fail-safe  $N$  by direct computation and calculated the ratio of fail-safe  $N$  to the number of published studies; and 2) followed up with re-analyses stratified by sample size and date of publication.

## Results

### Male Studies

Our search identified 20 studies of exclusively male samples or mixed sex samples for which results were available in male subjects separately (Table 1). These studies included 11,064 subjects (Table 2). The meta-analysis showed MAOA genotype to moderate an association of early life adversities (maltreatment plus other adversities) with later aggressive and antisocial outcomes across all male cohorts ( $p = .0044$ ) (Figure 1). This effect persisted: 1) on removal of each study individually ( $2.5 \times 10^{-6} < p < .018$ ) (Table 2); and 2) with iterative substitution of individual outcome variables for composite measures or averaged  $p$  values ( $6.6 \times 10^{-5} < p < .014$ ). Additional analyses showed the interaction significant in all but 1 of 1000 random combinations of study-specific dependent variables and environmental moderators ( $7.1 \times 10^{-6} < p < .06$ ). Results were significant for outcomes indexed to either childhood/adolescence ( $p = .032$ ) or adulthood ( $p = .008$ ); outcomes of aggression/violence ( $p = 3.9 \times 10^{-5}$ ), nonviolent antisocial behaviors ( $p = 7.2 \times 10^{-4}$ ), or combined indices ( $p = .022$ ); for dependent measures of continuous ( $p = .008$ ) but not dichotomous distribution ( $p = .36$ ); and for studies of both cross-sectional samples ( $p < .0045$ ) and longitudinally studied cohorts ( $p = .019$ ). Findings were unaffected by deletion of two nonwhite samples (19,24) ( $p = .0044$ ) or inclusion of four additional cohorts ( $n = 806$ ) of majority-male, mixed-sex samples (17,33,34) ( $p = .0034$ ).

**Stratified Analyses.** Low-activity MAOA genotype heightened risk for antisocial behavior among individuals exposed to maltreatment specifically ( $p = .00000082$ ) but not in tests of other childhood adversities ( $p = .40$ ). The interaction with maltreatment remained highly significant: 1) when deleting each study individually ( $2.2 \times 10^{-5} < p < 2.7 \times 10^{-7}$ ) (Table 3); 2) with iterative substitution of individual outcome measures for composite indices or mean  $p$  values ( $3.5 \times 10^{-5} < p < 1.4 \times 10^{-7}$ ); and 3) across all of 1000 random combinations of individual outcome measures ( $1.8 \times 10^{-9} < p < .005$ ). Among maltreatment studies, findings were again significant when analyzed separately for child/adolescent ( $p = 3.6 \times 10^{-5}$ ) and adult outcomes ( $p = .05$ ) and for outcomes of aggression/violence ( $p = .01$ ), nonviolent antisocial behaviors ( $p = 4.0 \times 10^{-4}$ ), and combined indices ( $p = 3.6 \times 10^{-6}$ ). They were significant also for dependent variables of both continuous ( $p = 1.4 \times 10^{-6}$ ) and dichotomous distribution ( $p = .02$ ); in cross-sectional studies ( $p = .01$ ) and studies of longitudinal cohorts ( $p = 3.5 \times 10^{-5}$ ); and where assessment of maltreatment exposure rested on family-based report only ( $p = 7.8 \times 10^{-7}$ ) or included nonfamilial informant sources ( $p = .022$ ). Here, too, results were unaltered by removal of two nonwhite cohorts (19,24) ( $p = 5.4 \times 10^{-7}$ ) or addition of the four majority-male mixed-sex samples (17,33,34) ( $p = 5.7 \times 10^{-7}$ ).

### Female Studies

We identified 12 studies involving female subjects only or separately analyzed female cohorts, with a total of 7588 subjects (Table 1). The meta-analysis showed no significant interaction of MAOA genotype with early life adversities (maltreatment and other adversities) across all studies ( $p = .77$ ).

**Stratified Analyses.** When analyzed separately, MAOA genotype predicted antisocial outcomes in interaction with childhood maltreatment ( $p = .020$ ) but not on exposure to other early adversities ( $p = .32$ ). Unlike male subjects, the interaction with maltreatment reflected an increased risk linked to high-activity MAOA genotype. On deletion of each study individually, however,

**Table 1.** Description of MAOA, Early Life Adversity, and Antisocial Behavior Studies Included in the Meta-Analysis

Source, Year	n	Male %	Race	Study Design	Moderator			Outcome			One-Tailed p Value <sup>b</sup>	
					Adversity Measure	Informant	Age	Antisocial Behavior	Informant	Age		Finding <sup>a</sup>
Caspi <i>et al.</i> (1) 2002	442	100%	White	Longitudinal	Maltreatment	INT (C), OBS, PR	3-11	CD, violent conviction, APD sx, violent disposition	INF, INT (C), OR, SR	11-18	Positive	.0050
Foley <i>et al.</i> (12) 2004	229	0%						CD	INT (C)		Negative	.1285 <sup>c</sup>
Haberstick <i>et al.</i> (13) 2005	514	100%	White	Longitudinal	Maltreatment	INT (C, P)	<18	CD	INT (C, P)	<18	Positive	.0200
Huizinga <i>et al.</i> (14) 2006	772	100%	White	Longitudinal	Maltreatment	INT (C), SR	<18	Conduct problems, violent convictions	INT, OR	16, 17, 22	Negative	.1423 <sup>d</sup>
Huizinga <i>et al.</i> (14) 2006	277	100%	White	Longitudinal	Maltreatment	SR	<17	CD, violent conviction, APD sx, violent disposition	INF, INT (C), OR, SR	14-28	Negative (opposite)	.7794 <sup>d</sup>
Kim-Cohen <i>et al.</i> (15) 2006	975	100%	White	Longitudinal	Maltreatment	INT (P)	≤7	ASB; attention/hyperactivity; emotional problems	PR, TR	7	Positive	.0145
Nilsson <i>et al.</i> (16) 2006	79	100%	White	Cross-sectional	Maltreatment; type of residence	INT (C), OR	≤16 or 19	Stealing, vandalism, violence	INT (C)	16, 19	Positive	.0078
Widom and Brzustowicz (17) 2006	261	67%	White	Longitudinal (case vs. control)	Maltreatment	OR	<12	CD sx, APD sx, violent behaviors	INT (C), OR, SR	<18-18+	Positive	.0143 <sup>d</sup>
Frazzetto <i>et al.</i> (18) 2007	148	62%	NW								Negative	.5000 <sup>d</sup>
Frazzetto <i>et al.</i> (18) 2007	82	100%	White	Cross-sectional (case vs. control)	Early adverse experiences	SR	<16	Physical aggression	SR	m = 30.88	Positive	.0020
Sjoberg <i>et al.</i> (26) 2007	153	0%									Negative	.4230
Sjoberg <i>et al.</i> (26) 2007	117	0%	White	Cross-sectional	Maltreatment; type of residence	SR	≤16 or 19	Stealing, vandalism, violence	INT (C)	16-19	Negative (opposite)	.9082
Vanyukov <i>et al.</i> (27) 2007	144	100%	White	Longitudinal (high risk)	Parental involvement	SR	12-18	CD, ADHD	INT (C, P)	12-18	Partial opposite	.6517 <sup>d</sup>
Ducci <i>et al.</i> (37) 2008	187	0%	White	Cross-sectional (case vs. control)	Maltreatment	INT (C), OR	<16	APD sx	INT (C)	m = 37.80	Positive	.0002
Hart and Marmorstein (28) 2009	672	100%	NW UNR	Longitudinal	Neighborhood characteristics	OR	r = 11-21	Aggression	SR	r = 11-23	Positive	.0250
Prom-Wormley <i>et al.</i> (36) 2009	721	0%	White	Longitudinal	Maltreatment	PR, SR	<18	CD	PR, SR	<18	Opposite	.9750
van der Vegt <i>et al.</i> (19) 2009	239	100%	NW	Cross-sectional	Maltreatment	INF	≤15	Externalizing behaviors (aggression, delinquency)	PR	<15	Negative (opposite)	.9000
Weder <i>et al.</i> (33) 2009	114	66%	NW	Cross-sectional (case vs. control)	Maltreatment	INT (P, C), SR, OR	≤15	Aggression, rule breaking, inattention	TR	≤15	Partial positive	.0359 <sup>d</sup>
Beach <i>et al.</i> (20) 2010	244	100%	White	Longitudinal	Maltreatment	INT (C)	<18	APD sx	INT (C)	m = 46.48	Negative	.0300
Beach <i>et al.</i> (20) 2010	294	0%								m = 44.95	Positive	.0240
Beaver <i>et al.</i> (29) 2010	420	100%	White	Longitudinal	Protective-risk index	SR	<18	Incarceration, anger/hostility	OR, SR	r = 24-32	Partial positive	.0694 <sup>d</sup>
Beaver <i>et al.</i> (29) 2010	493	0%									Negative	.1539 <sup>d</sup>

Table 1. (continued)

Source, Year	n	Male %	Race	Study Design	Moderator			Outcome				One-Tailed p Value <sup>b</sup>
					Adversity Measure	Informant	Age	Antisocial Behavior	Informant	Age	Finding <sup>a</sup>	
Derringer <i>et al.</i> (21) 2010	595	100%	White	Longitudinal	Maltreatment	INT (C), SR	<18	CD sx, APD sx	INT (C)	17, 21, 25	Negative	.3520 <sup>d</sup>
	246	0%									Partial positive	.1057 <sup>d</sup>
Edwards <i>et al.</i> (22) 2010	186	100%	White	Longitudinal (high risk)	Maltreatment	SR	<6	Externalizing behaviors (aggression, delinquency)	PR, SR, TR	6–22	Negative	.0675
Enoch <i>et al.</i> (30) 2010	3182	100%	White	Longitudinal	Family adversity, stressful life events	PR	PN to 7	Conduct problems, ADHD	PR	4, 7	Negative	.3936 <sup>d</sup>
	3976	0%									Partial positive	.1635 <sup>d</sup>
Wakschlag <i>et al.</i> (31) 2010	78	100%	White	Longitudinal (high risk)	Prenatal smoking	DT, PR	PN	CD sx	PR	m = 15	Positive	.0160 <sup>d</sup>
Aslund <i>et al.</i> (23) 2011	99	0%									Opposite	.9990 <sup>d</sup>
	780	100%	White	Cross-sectional	Maltreatment	INT (C)	<18	Stealing, vandalism, violence	SR	r = 17–18	Positive	.0048 <sup>d</sup>
Lee (32) 2011	882	0%									Opposite	.9995 <sup>d</sup>
	672	100%	White	Longitudinal	Deviant peer behavior	SR	m = 15.65	Overt/covert ASB	SR	mr = 15–22	Partial opposite	.9641 <sup>d</sup>
Reti <i>et al.</i> (34) 2011	283	52%	White	Cross-sectional	Maltreatment	INT (C)	<18	APD sx	INT (C)	18+	Opposite	.9236
Cicchetti <i>et al.</i> (24) 2012	312	100%	NW	Cross-sectional (high risk)	Maltreatment	OR	≤12	CD sx, ASB, externalizing behaviors (aggression, delinquency)	INF, INT (C), TR	≤12	Partial positive	.1190 <sup>d</sup>
Fergusson <i>et al.</i> (25) 2012	351	100%	White	Longitudinal	Maltreatment; PN smoking; material deprivation	INT (C), SR	PN to 15	Violent/property offenses, conduct problems, hostility	INT (C), SR, OR	15–21; 25, 30	Positive	.0192 <sup>d</sup>
McGrath <i>et al.</i> (38) 2012	192	0%	White	Cross-sectional (high risk)	Maltreatment	SR	<18	Conduct problems, impulsive-sensation seeking, interpersonal aggression, PM smoking	SR	m = 42.9	Negative (opposite)	.9066

ADHD, attention-deficit/hyperactivity disorder; APD, antisocial personality disorder; ASB, antisocial behaviors; C, child; CD, conduct disorder; DT, drug test; INF, other informant; INT, interview; m, mean age; mr, mean age range; NW, nonwhite; OBS, observation; OR, official record; P, parent; PN, prenatal; PR, parent-report; r, age range; SR, self-report; sx, symptoms; TR, teacher-report; UNS, unspecified.

<sup>a</sup>Positive indicates a significant ( $p < .05$ ) interaction effect with the low-activity *MAOA* variant, as presented in the original study report, Negative indicates no interaction effect ( $p > .05$ ), and Opposite indicates a significant ( $p < .05$ ) interaction with the high-activity *MAOA* variant. Partial indicates a significant interaction effect for one or more, but not all, measures tested in a study with multiple dependent variables.

<sup>b</sup>One-tailed  $p$  value, with smaller values ( $p < .50$ ) indicating greater sensitivity among low-activity *MAOA* subjects and larger values ( $p > .50$ ) indicating greater sensitivity among high-activity *MAOA* subjects.

<sup>c</sup>From footnoted observations that were not a component of primary analyses in Caspi *et al.* (1) 2002.

<sup>d</sup>Denotes a weighted mean  $p$  value created when multiple dependent variables and/or multiple environmental moderators were analyzed separately in the published report.

**Table 2.** Studies Included in the All Male Group Meta-Analysis

Source, Year	Total Number of Participants	One-Tailed <i>p</i> Value	Fisher <i>p</i> Value after Study Exclusion
Caspi <i>et al.</i> (1) 2002	442	.0050	$1.02 \times 10^{-2}$
Foley <i>et al.</i> (12) 2004	514	.0200	$9.32 \times 10^{-3}$
Haberstick <i>et al.</i> (13) 2005	772	.1423	$7.15 \times 10^{-3}$
Huizinga <i>et al.</i> (14) 2006	277	.7794	$3.50 \times 10^{-3}$
Kim-Cohen <i>et al.</i> (15) 2006	975	.0145	$1.73 \times 10^{-2}$
Nilsson <i>et al.</i> (16) 2006	79	.0078	$5.11 \times 10^{-3}$
Frazzetto <i>et al.</i> (18) 2007	82	.0020	$5.27 \times 10^{-3}$
Vanyukov <i>et al.</i> (27) 2007	144	.6517	$4.10 \times 10^{-3}$
Hart and Marmorstein (28) 2009	672	.0250	$1.08 \times 10^{-2}$
van der Vegt <i>et al.</i> (19) 2009	239	.9000	$3.28 \times 10^{-3}$
Beach <i>et al.</i> (20) 2010	244	.0300	$6.14 \times 10^{-3}$
Beaver <i>et al.</i> (29) 2010	420	.0694	$6.73 \times 10^{-3}$
Derringer <i>et al.</i> (21) 2010	595	.3520	$4.65 \times 10^{-3}$
Edwards <i>et al.</i> (22) 2010	186	.0675	$5.44 \times 10^{-3}$
Enoch <i>et al.</i> (30) 2010	3182	.3936	$2.48 \times 10^{-6}$
Wakschlag <i>et al.</i> (31) 2010	78	.0160	$4.95 \times 10^{-3}$
Aslund <i>et al.</i> (23) 2011	780	.0048	$1.78 \times 10^{-2}$
Lee (32) 2011	672	.9641	$1.28 \times 10^{-3}$
Cicchetti <i>et al.</i> (24) 2012	312	.1190	$5.61 \times 10^{-3}$
Fergusson <i>et al.</i> (25) 2012	399	.0192	$8.05 \times 10^{-3}$
Total	11,064		
Average Sample Size	553		.0044

this finding lost significance with removal of either of two study cohorts ( $.004 < p < .97$ ) (Table 4).


### Publication Bias

Our results corroborate the sentinel observation of Caspi *et al.* (1) that childhood maltreatment predicts antisocial outcomes more strongly in male subjects of low-activity, compared with high-activity, *MAOA* genotype, and do not show this interaction extended to other categories of early life adversity or to female subjects. To render the *MAOA* × maltreatment interaction in male subjects nonsignificant ( $p > .05$ ) would require > 93 unpublished analyses or undiscovered studies of null effect ( $p = .50$ ) and equal average sample size ( $n = 447$ ). This yields a fail-safe ratio of seven studies not included for each maltreatment study of male subjects included in the meta-analysis. The *MAOA* × maltreatment interaction also proved significant in analyses restricted to: 1) studies with samples either larger ( $p = 1.7 \times 10^{-4}$ ) or smaller ( $p = .01$ ) than Caspi *et al.* (1); and 2) either recent (dated 2010–2012;  $p = 3.0 \times 10^{-4}$ ) or early replication attempts (2004–2009;  $p = .005$ ).

### Discussion

In their provocative first study of G×E interaction, Caspi *et al.* (1) reported that common polymorphic variation in *MAOA* moderated the influence of childhood maltreatment on boys' later aggressive and antisocial behaviors, as seen in a longitudinally studied, normal population. Our purpose was to determine whether this finding replicated in subsequent research addressed to the same hypothesis, when again examined in primarily nonclinical samples and extended to studies of other early life adversities or to female subjects. Across male cohorts, the meta-analysis showed a moderately reliable interaction of *MAOA*

variation and environmental risk factors, with childhood adversities presaging antisocial outcomes more strongly in persons of low-activity, compared with high-activity, *MAOA* genotype [as in Caspi *et al.* (1)]. Moreover, analyses stratified by category of early life adversity showed this finding was accounted for principally by the interaction of *MAOA* and childhood maltreatment ( $p = 2.0 \times 10^{-7}$ ) and therefore, where environmental risk most closely matched the sentinel study (1). The interaction with childhood maltreatment also proved robust to sensitivity analyses and generalized across studies of either cross-sectional or longitudinal design and studies in which maltreatment exposure was assessed by family (self, parent) report only or included independent informant sources. It is noteworthy, too, that *MAOA* variation interacted with childhood maltreatment to predict outcomes referenced to both childhood/adolescence and adulthood, dependent measures of both continuous and categorical distribution, and both violent and non-violent antisocial behaviors. The latter finding suggests that the low-activity *MAOA* genotype heightens maltreatment-dependent



**Figure 1.** Forest plot of 20 male samples for the interaction of *MAOA* genotype and early life adversities on aggressive and antisocial behavior. Icons indicate the one-tailed *p* value for each sample, where lower values denote a greater sensitivity to adversity with low-activity *MAOA* genotype and high values denote a greater sensitivity with high-activity *MAOA* genotype. The size of the icon reflects relative sample size. Squares mark studies that indexed adversity specifically to childhood maltreatment; circles indicate studies of other childhood adversities; and diamonds indicate studies that included both maltreatment and other childhood adversities. The red triangle depicts the overall result of the meta-analysis for all-male samples (two-tailed). Studies marked with an asterisk were included in the prior meta-analysis by Taylor and Kim-Cohen (11).

**Table 3.** Studies Included in the All Male, Maltreatment Group Meta-Analysis

Source, Year	Total Number of Participants	One-Tailed <i>p</i> Value	Fisher <i>p</i> Value after Study Exclusion
Caspi <i>et al.</i> (1) 2002	442	.0050	$8.59 \times 10^{-6}$
Foley <i>et al.</i> (12) 2004	514	.0200	$5.63 \times 10^{-6}$
Haberstick <i>et al.</i> (13) 2005	772	.1423	$8.20 \times 10^{-7}$
Huizinga <i>et al.</i> (14) 2006	277	.7794	$3.40 \times 10^{-7}$
Kim-Cohen <i>et al.</i> (15) 2006	975	.0145	$8.59 \times 10^{-6}$
Nilsson <i>et al.</i> (16) 2006	79	.0183	$1.23 \times 10^{-6}$
van der Vegt <i>et al.</i> (19) 2009	239	.9000	$2.70 \times 10^{-7}$
Beach <i>et al.</i> (20) 2010	244	.0300	$2.25 \times 10^{-6}$
Derringer <i>et al.</i> (21) 2010	595	.3520	$3.80 \times 10^{-7}$
Edwards <i>et al.</i> (22) 2010	186	.0675	$1.51 \times 10^{-6}$
Aslund <i>et al.</i> (23) 2011	780	.0048	$2.24 \times 10^{-5}$
Cicchetti <i>et al.</i> (24) 2012	312	.1190	$1.59 \times 10^{-6}$
Fergusson <i>et al.</i> (25) 2012	399	.0051	$7.46 \times 10^{-6}$
Total	5814		
Average Sample Size	447		.00000082

risk for a range of conduct problems and not aggression or criminal violence specifically.

These findings are consistent with a broader literature on early risk factors for antisocial behavior, in which maltreatment indicators like domestic violence, physical abuse, neglect, and parental rejection figure prominently (52–60). They also accord with findings of a recent twin study, in which maltreatment increased children's risk for conduct problems as a function of latent genetic risk, defined by twin-pair zygosity and co-twin diagnostic status for conduct disorder (61). Although biological mechanisms underlying these associations remain unknown, environmental insults like maltreatment presumably compound or otherwise interact with neurobehavioral correlates of *MAOA* to augment aggressive or antisocial potential. Relatedly, individuals of low-activity, compared with high-activity, *MAOA* genotype have performed more poorly on some executive processing tasks, such as tests of working memory and attentional control, and exhibited reduced task-dependent activations of frontal brain regions supporting these processes (62–66). The low-activity *MAOA* genotype has been linked as well to altered neural responses to affective stimuli, including enhanced amygdala reactions to facial expressions of emotion or emotion recall, lesser engagement of prefrontal regulatory regions, and disrupted functional and effective (top-down) connectivity within cortico-lymbic circuitry of emotion processing (63,67–69). It is possible that early maltreatment either exacerbates these neural deficits or engenders antagonistic and antisocial motivations that are abetted by *MAOA*-modulated impairments in inhibitory control.

In contrast to studies of childhood maltreatment, *MAOA* genotype did not moderate the aggregate effects of other early life adversities in male subjects. The collection of environmental risks sampled in these investigations was quite variable, however, and it may be premature to conclude that *MAOA* interacts only with maltreatment to affect antisocial outcomes. For instance, risk associated with maternal prenatal smoking varied by *MAOA* genotype in each of the two studies that examined this variable (weighted  $p = .024$ ) (25,31). We should note that all but one attempted replication included in the prior meta-analysis by Taylor and Kim-Cohen (11) were categorized here as studies of maltreatment. The one such investigation that we placed instead

in the category of other adversities used a measure of family relationships that, although encompassing abuse, emphasized a broader range of difficulties (e.g., separations, disability of self or sibling, marital problems, and parental psychopathology) (18). Nonetheless, this study found the interaction of *MAOA* genotype and family difficulties to also predict male aggression, so that if included in meta-analysis with studies focusing more explicitly on maltreatment, the outcome does not differ ( $p = 4.3 \times 10^{-7}$ ).

Unlike male subjects, *MAOA* variation did not interact with early life adversities across 12 female cohorts, but like male subjects, the interaction was significant in studies of childhood maltreatment alone. Yet, among girls who were maltreated, the high-activity, not low-activity, *MAOA* genotype was more strongly associated with antisocial outcomes, although the interaction was weak and lost significance with deletion of some individual studies. Still, it is interesting that a combination of high-activity *MAOA* genotype and environmental risk predicted delinquent behavior in several studies of adolescent girls and across different groups of investigators (23,31,36,38). Elsewhere, symptoms of dysthymia have been found greater in women who were maltreated in childhood and carry the *MAOA* high-activity (4-repeat) variant than among women of homozygous, low-activity genotype (70). At present, it is unclear how a reversal of allelic association between male and female subjects might be explained, and existing literature provides few clues. Incomplete X-inactivation at the *MAOA* locus could conceivably produce a different expression profile in women, possibly yielding a sex difference in *MAOA* product (50,51). Also potentially relevant to *MAOA* expression is some evidence that CpG residues in the *MAOA* promoter are hypermethylated in women, compared with men, and that differential methylation may be greatest among women of low-activity *MAOA* genotype (71). A third possibility is that *MAOA* interacts with sex differences in perinatal androgen exposure to affect brain development or that gonadal hormones modulate genotype-dependent variation in *MAOA* expression in adolescence (26,70). These suggestions are highly speculative, however, and do not yet provide a clear mechanism for a bidirectional association of *MAOA* genotype with antisocial behavior among maltreated cohorts of different sexes. Nonetheless, additional research may be warranted to determine if the findings suggesting heightened susceptibility for the high-activity genotype in female subjects emerge more prominently in a larger literature.

To recapitulate, our meta-analysis confirms the first seminal study of G×E interaction reported by Caspi *et al.* (1). We find that

**Table 4.** Studies Included in the All Female, Maltreatment Group Meta-Analysis

Source, Year	Total Number of Participants	One-Tailed <i>p</i> Value	Fisher <i>p</i> Value after Study Exclusion
Caspi <i>et al.</i> (1) 2002	229	.1285	$9.88 \times 10^{-3}$
Sjoberg <i>et al.</i> (26) 2007	117	.9494	$2.85 \times 10^{-2}$
Ducci <i>et al.</i> (37) 2008	187	.0002	$3.73 \times 10^{-3}$
Prom-Wormley <i>et al.</i> (36) 2009	721	.9750	$1.39 \times 10^{-1}$
Beach <i>et al.</i> (20) 2010	294	.0240	$3.98 \times 10^{-3}$
Derringer <i>et al.</i> (21) 2010	246	.1057	$8.54 \times 10^{-3}$
Aslund <i>et al.</i> (23) 2011	882	.9995	$9.68 \times 10^{-1}$
McGrath <i>et al.</i> (38) 2012	192	.9066	$3.08 \times 10^{-2}$
Total	2868		
Average Sample Size	359		.02

MAOA variation moderates effects of early life adversity on male subjects' aggressive and antisocial behaviors, and this interaction is attributable to studies that, like the initial report, delimit adversity to experiences of maltreatment, such as physical and sexual abuse, harsh discipline, neglect, assault, or other ill-treatment. Significance of the weighted Z score for direct replications (i.e., maltreatment studies) is substantial ( $p = 8.2 \times 10^{-7}$ ) and supported by a sizable fail-safe ratio. Of course, even a positive meta-analysis does not exhaust validity challenges or vouchsafe a true association. For instance, publication bias may be indicated if reported replications aggregate among smaller rather than larger, better powered studies or in early, but not later, studies. Here, however, we found the interaction of MAOA genotype and childhood maltreatment no less likely to replicate in studies with sample sizes larger (or smaller) than Caspi *et al.* (1) or in studies published later (in the last 3 years) or before 2010. Indeed, the interaction also proves significant ( $p = 5.1 \times 10^{-4}$ ) if restricted only to maltreatment studies not included in the preceding (positive) meta-analysis of Taylor and Kim-Cohen (11). Finally, considering the recent, parallel meta-analysis of 5-HTTLPR variation, life stress, and depression (7) alongside our study suggests that the two novel investigations spawning these literatures (1,2) reflect not only prominent but also durable examples of G×E interaction. Both analyses also highlight points of methodology affecting study outcomes, such as type of early adversity (here) or in studies of 5-HTTLPR and depression, stressor type and quality of measurement, which may usefully guide future research aimed at elucidation of etiologic mechanisms. Whether these two instances reflect a more general role of G×E interactions in the genesis of major psychopathologies is unknown, however, and awaits the emergence of comparable literatures addressed to other disorders, genes, and environmental risk factors.

*This work was supported in part by National Institutes of Health Grant PO1 HL040962 (SBM) and National Science Foundation Integrative Graduate Education and Research Traineeship Program 0549352 (ALB).*

*The authors report no biomedical financial interests or potential conflicts of interest.*

*Editor's Note: Minor editorial corrections and clarifications were incorporated into this final version of the article since the article-in-press version was made available online.*

- Caspi A, McClay J, Moffitt TE, Mill J, Martin J, Craig IW, *et al.* (2002): Role of genotype in the cycle of violence in maltreated children. *Science* 297:851–854.
- Caspi A, Sugden K, Moffitt TE, Taylor A, Craig IW, Harrington H, *et al.* (2003): Influence of life stress on depression: Moderation by a polymorphism in the 5-HTT gene. *Science* 301:386–389.
- Munafò MR, Durrant C, Lewis G, Flint J (2009): Gene × environment interactions at the serotonin transporter locus. *Biol Psychiatry* 65:211–219.
- Risch N, Herrell R, Lehner T, Liang KY, Eaves L, Hoh J, *et al.* (2009): Interaction between the serotonin transporter gene (5-HTTLPR), stressful life events, and risk of depression: A meta-analysis. *JAMA* 301:2462–2471.
- Rutter M, Thapar A, Pickles A (2009): Gene-environment interactions: Biologically valid pathway or artifact? *Arch Gen Psychiatry* 66:1287–1289.
- Uher R, McGuffin P (2010): The moderation by the serotonin transporter gene of environmental adversity in the etiology of depression: 2009 update. *Mol Psychiatry* 15:18–22.
- Karg K, Burmeister M, Shedden K, Sen S (2011): The serotonin transporter promoter variant (5-HTTLPR), stress, and depression meta-analysis revisited: Evidence of genetic moderation. *Arch Gen Psychiatry* 68:444–454.
- Sabol SZ, Hu S, Hamer D (1998): A functional polymorphism in the monoamine oxidase A gene promoter. *Hum Genet* 103:273–279.
- Denney RM, Koch H, Craig IW (1999): Association between monoamine oxidase A activity in human male skin fibroblasts and genotype of the MAOA promoter-associated variable number tandem repeat. *Hum Genet* 195:542–551.
- Deakert J, Catalano M, Sygailo YV, Bosi M, Okladnova O, Di Bella D, *et al.* (1999): Excess of high activity monoamine oxidase A gene promoter alleles in females patients with panic disorder. *Hum Mol Genet* 8:621–624.
- Taylor A, Kim-Cohen J (2007): Meta-analysis of gene-environment interactions in developmental psychopathology. *Dev Psychopathol* 19:1029–1037.
- Foley DL, Eaves LJ, Wormley B, Silberg JL, Maes HH, Kuhn J, Riley B (2004): Childhood adversity, monoamine oxidase A genotype, and risk for conduct disorder. *Arch Gen Psychiatry* 61:738–744.
- Haberstick BC, Lessem JM, Hopfer CJ, Smolen A, Ehringer MA, Timberlake D, Hewitt JK (2005): Monoamine oxidase A (MAOA) and antisocial behaviors in the presence of childhood and adolescent maltreatment. *Am J Med Genet B Neuropsychiatr Genet* 135:59–64.
- Huizinga D, Haberstick BC, Smolen A, Menard S, Young SE, Corley RP, *et al.* (2006): Childhood maltreatment, subsequent antisocial behavior, and the role of monoamine oxidase A genotype. *Biol Psychiatry* 60:677–683.
- Kim-Cohen J, Caspi A, Taylor A, Williams B, Newcombe R, Craig IW, Moffitt TE (2006): MAOA, maltreatment, and gene-environment interaction predicting children's mental health: New evidence and a meta-analysis. *Mol Psychiatry* 11:903–913.
- Nilsson KW, Sjöberg RL, Damberg M, Leppert J, Ohrvik J, Alm PO, *et al.* (2006): Role of monoamine oxidase A genotype and psychosocial factors in male adolescent criminal activity. *Biol Psychiatry* 59:121–127.
- Widom CS, Brzustowicz LM (2006): MAOA and the "Cycle of Violence": Childhood abuse and neglect, MAOA genotype, and risk for violent and antisocial behavior. *Biol Psychiatry* 60:684–689.
- Frazzetto G, Di Lorenzo G, Carola V, Proietti L, Sokolowska E, Siracusano A, *et al.* (2007): Early trauma and increased risk for physical aggression during adulthood: The moderating role of MAOA genotype. *PLoS One* 2:e486.
- van der Vegt EJM, Oostra BA, Arias-Vásquez A, van der Ende J, Verhulst FC, Tiemeier H (2009): High activity of monoamine oxidase A is associated with externalizing behaviour in maltreated and nonmaltreated adoptees. *Psychiatr Genet* 19:209–211.
- Beach SRH, Brody GH, Gunter TD, Packer H, Wernett P, Philibert RA (2010): Child maltreatment moderates the association of MAOA with symptoms of depression and antisocial personality disorder. *J Fam Psychol* 24:12–20.
- Derringer J, Krueger RF, Irons D, Iacono WG (2010): Harsh discipline, childhood sexual assault, and MAOA genotype: An investigation of main and interactive effects on diverse clinical externalizing outcomes. *Behav Genet* 40:639–648.
- Edwards AC, Dodge KA, Latendresse SJ, Lansford JE, Bates JE, Pettit GS, *et al.* (2010): MAOA-uVNTR and early physical discipline interact to influence delinquent behavior. *J Child Psychol Psychiatry* 51:679–687.
- Åslund C, Nordquist N, Comasco E, Leppert J, Orelund L, Nilsson K (2011): Maltreatment, MAOA, and delinquency: Sex differences in gene-environment interaction in a large population-based cohort of adolescents. *Behav Genet* 41:262–272.
- Cicchetti D, Rogosch FA, Thibodeau EL (2012): The effects of child maltreatment on early signs of antisocial behavior: Genetic moderation by tryptophan hydroxylase, serotonin transporter, and monoamine oxidase A genes. *Dev Psychopathol* 24:907–928.
- Fergusson DM, Boden JM, Horwood LJ, Miller A, Kennedy MA (2012): Moderating role of the MAOA genotype in antisocial behaviour. *Br J Psychiatry* 200:116–123.
- Sjöberg RL, Nilsson KW, Wargelius HL, Leppert J, Lindström L, Orelund L (2007): Adolescent girls and criminal activity: Role of MAOA-LPR genotype and psychosocial factors. *Am J Med Genet B Neuropsychiatr Genet* 144B:159–164.
- Vanyukov MM, Maher BS, Devlin B, Kirillova GP, Kirisci L, Yu LM, Ferrell RE (2007): The MAOA promoter polymorphism, disruptive behavior disorders, and early onset substance use disorder: Gene-environment interaction. *Psychiatr Genet* 17:323–332.
- Hart D, Marmorstein NR (2009): Neighborhoods and genes and everything in between: Understanding adolescent aggression in social and biological contexts. *Dev Psychopathol* 21:961–973.



29. Beaver KM, DeLisi M, Vaughn MG, Barnes JC (2010): Monoamine oxidase A genotype is associated with gang membership and weapon use. *Compr Psychiatry* 51:130–134.
30. Enoch MA, Steer CD, Newman TK, Gibson N, Goldman D (2010): Early life stress, MAOA, and gene-environment interactions predict behavioral disinhibition in children. *Genes Brain Behav* 9:65–74.
31. Wakschlag LS, Kistner EO, Pine DS, Biesecker G, Pickett KE, Skol AD, et al. (2010): Interaction of prenatal exposure to cigarettes and MAOA genotype in pathways to youth antisocial behavior. *Mol Psychiatry* 15:928–937.
32. Lee SS (2011): Deviant peer affiliation and antisocial behavior: Interaction with monoamine oxidase A (MAOA) genotype. *J Abnorm Child Psychol* 39:321–332.
33. Weder N, Yang B, Douglas-Palumberi H, Massey J, Krystal JH, Gelernter J, Kaufman J (2009): MAOA genotype, maltreatment, and aggressive behavior: The changing impact of genotype at varying levels of trauma. *Biol Psychiatry* 65:417–424.
34. Reti IM, Xu JZ, Yanofski J, McKibben J, Uhart M, Cheng YJ, et al. (2011): Monoamine oxidase A regulates antisocial personality in whites with no history of physical abuse. *Compr Psychiatry* 52:188–194.
35. Sjöberg RL, Ducci F, Barr CS, Newman TK, Dell'Osso L, Virkkunen M, Goldman D (2008): A non-additive interaction of a functional MAO-A VNTR and testosterone predicts antisocial behavior. *Neuropsychopharmacology* 33:425–430.
36. Prom-Wormley EC, Eaves LJ, Foley DL, Gardner CO, Archer KJ, Wormley BK, et al. (2009): Monoamine oxidase A and childhood adversity as risk factors for conduct disorder in females. *Psychol Med* 39:579–590.
37. Ducci F, Enoch M, Hodgkinson C, Xu K, Catena M, Robin RW, Goldman D (2008): Interaction between a functional MAOA locus and childhood sexual abuse predicts alcoholism and antisocial personality disorder in adult women. *Mol Psychiatry* 13:334–347.
38. McGrath L, Mustanski B, Metzger A, Pine D, Kistner-Griffin E, Cook E, Wakschlag LS (2012): A latent modeling approach to genotype-phenotype relationships: Maternal problem behavior clusters, prenatal smoking, and MAOA genotype. *Arch Womens Ment Health* 15:269–282.
39. Whitlock MC (2005): Combining probability from independent tests: The weighted Z-method is superior to Fisher's approach. *J Evol Biol* 18:1368–1373.
40. Hedges LV, Olkin I (1985): *Statistical Methods for Meta-Analysis*. New York: Academic.
41. Richards JB, Waterworth D, O'Rahilly S, Hivert MF, Loos RJ, Perry JR, et al. (2009): A genome-wide association study reveals variants in ARL15 that influence adiponectin levels. *PLoS Genet* 5:e1000768.
42. Hwang D, Rust AG, Ramsey S, Smith JJ, Leslie DM, Weston AD, et al. (2005): A data integration methodology for systems biology. *Proc Natl Acad Sci U S A* 102:17296–17301.
43. Young SE, Smolen A, Hewitt JK, Haberstick BC, Stallings MC, Corley RP, Crowley TJ (2006): Interaction between MAO-A genotype and maltreatment in the risk for conduct disorder: Failure to confirm in adolescent patients. *Am J Psychiatry* 163:1019–1025.
44. Reif A, Rosler M, Freitag CM, Schneider M, Eujen A, Kissling C, et al. (2007): Nature and nurture predispose to violent behavior: Serotonergic genes and adverse childhood environment. *Neuropsychopharmacology* 32:2375–2383.
45. Tikkanen R, Ducci F, Goldman D, Holi M, Lindberg N, Tiihonen J, Virkkunen M (2010): MAOA alters the effects of heavy drinking and childhood physical abuse on risk for severe impulsive acts of violence among alcoholic violent offenders. *Alcohol Clin Exp Res* 34:853–860.
46. Huang YY, Cate SP, Battistuzzi C, Oquendo MA, Brent D, Mann JJ (2004): An association between a functional polymorphism in the monoamine oxidase a gene promoter, impulsive traits and early abuse experiences. *Neuropsychopharmacology* 29:1498–1505.
47. Kinnally EL, Huang Y, Haverly R, Burke AK, Galfalvy H, Brent DP, et al. (2009): Parental care moderates the influence of MAOA-uVNTR genotype and childhood stressors on trait impulsivity and aggression in adult women. *Psychiatr Genet* 19:126–133.
48. Nilsson KW, Wargelius H-L, Sjöberg RL, Leppert J, Oreland L (2008): The MAO-A gene, platelet MAO-B activity and psychosocial environment in adolescent female alcohol-related problem behaviour. *Drug Alcohol Depend* 93:51–62.
49. Prichard Z, Mackinnon A, Jorm AF, Easteal S (2008): No evidence for interaction between MAOA and childhood adversity for antisocial behavior. *Am J Med Genet B Neuropsychiatr Genet* 147B:228–232.
50. Benjamin D, Van Badel I, Craig I (2000): A novel expression based approach for assessing the inactivation status of human X-linked genes. *Eur J Hum Genet* 8:103–108.
51. Carrel L, Willard H (2005): X-inactivation profile reveals extensive variability in X-linked gene expression in females. *Nature* 434:400–404.
52. Jaffee SR, Caspi A, Moffitt TE, Taylor A (2004): Physical maltreatment victim to antisocial child: Evidence of an environmentally mediated process. *J Abnorm Psychol* 113:44–55.
53. Fergusson DM, Horwood JL (1998): Exposure to interparental violence in childhood and psychosocial adjustment in young adulthood. *Child Abuse Negl* 22:339–357.
54. Jaffee SR, Moffitt TE, Caspi A, Taylor A, Arseneault L (2002): Influence of adult domestic violence on children's internalizing and externalizing problems: An environmentally informative twin study. *J Am Acad Child Adolesc Psychiatry* 41:1095–1103.
55. Caspi A, Moffitt TE, Morgan J, Rutter M, Taylor A, Arseneault L, et al. (2004): Maternal expressed emotion predicts children's antisocial behavior problems: Using monozygotic-twin differences to identify environmental effects on behavioral development. *Dev Psychol* 40:149–161.
56. Farrington DP, Hawkins JD (1991): Predicting participation, early onset and later persistence in officially recorded offending. *Crim Behav Mental Health* 1:1–33.
57. Luntz BK, Widom CS (1994): Antisocial personality disorder in abused and neglected children grown up. *Am J Psychiatry* 151:670–674.
58. Dodge KA, Sherrill MR (2007): The interaction of nature and nurture in antisocial behavior. In: Flannery DJ, Vazsonyi AT, Waldman ID, editors. *The Cambridge Handbook of Violent Behavior and Aggression*. New York: Cambridge University Press, 215–242.
59. Lansford JE, Dodge KA, Pettit GS, Bates JE, Crozier J, Kaplow J (2002): A 12-year prospective study of the long-term effects of early child physical maltreatment on psychological, behavioral, and academic problems in adolescence. *Arch Pediatr Adolesc Med* 156:824–830.
60. McCord J (1992): Questioning the value of punishment. *Social Problems* 38:167–176.
61. Jaffee SR, Caspi A, Moffitt TE, Dodge KA, Rutter M, Taylor A, Tully LA (2005): Nature x nurture: Genetic vulnerabilities interact with physical maltreatment to promote conduct problems. *Dev Psychopathol* 17:67–84.
62. Fan J, Fossella J, Sommer T, Wu Y, Posner MI (2003): Mapping the genetic variation of executive attention onto brain activity. *Proc Natl Acad Sci U S A* 100:7406–7411.
63. Meyer-Lindenberg A, Buckholtz JW, Kolachana B, Hariri A, Pezawas L, Blasi G, et al. (2006): Neural mechanisms of genetic risk for impulsivity and violence in humans. *Proc Natl Acad Sci U S A* 103:6269–6274.
64. Cerasa A, Gioia MC, Fera FP, L, Kiguori M, Lanza P, Muglia M, et al. (2008): Ventr-lateral prefrontal activity during working memory is modulated by MAO A genetic variation. *Brain Res* 1201:114–121.
65. Passamonti L, Fera F, Magariello A, Cerasa A, Gioia MC, Muglia M, et al. (2006): Monoamine oxidase-A genetic variations influence brain activity associated with inhibitory control: New insight into the neural correlates of impulsivity. *Biol Psychiatry* 59:334–340.
66. Enge S, Fleischhauer M, Lesch K-P, Reif A, Strobel A (2011): Serotonergic modulation in executive functioning: Linking genetic variations to working memory performance. *Neuropsychologia* 49:3776–3785.
67. Buckholtz JW, Meyer-Lindenberg A (2008): MAOA and the neuro-genetic architecture of human aggression. *Trends Neurosci* 31:120–129.
68. Buckholtz JW, Callicott JH, Kolachana B, Hariri AR, Goldberg TE, Genderson M, et al. (2008): Genetic variation in MAOA modulates ventromedial prefrontal circuitry mediating individual differences in human personality. *Mol Psychiatry* 13:313–324.
69. Lee B-T, Ham BJ (2008): Monoamine oxidase A-uVNTR genotype affects limbic brain activity in response to affective facial stimuli. *Neuroreport* 19:515–519.
70. Nikulina V, Widom CS, Brzustowicz LM (2012): Child abuse and neglect, MAOA, and mental health outcomes: A prospective examination. *Biol Psychiatry* 71:350–357.
71. Philibert RA, Gunter TD, Beach SRH, Brody GH, Madan A (2008): MAOA methylation is associated with nicotine and alcohol dependence in women. *Am J Med Genet B Neuropsychiatr Genet* 147B:565–570.